

PATIENT NAME:
DATE OF BIRTH:
DATE OF EXAM:
PHYSICIAN:

THYROID ULTRASOUND

TECHNIQUE

Utilizing real-time imaging, multiple gray scale sector images of the thyroid gland were obtained.

FINDINGS

The thyroid gland is in a normal cervical location. The bilateral thyroid lobes are heterogeneous. There is a 0.80 x 0.67 x 0.66 cm hypoechoic nodule with hyperechoic echoes and 0.94 x 0.28 cm hypoechoic nodule in the right lobe. There is a 0.87 x 0.87 x 0.63 cm hyperechoic nodule with calcification and 1.31 x 0.97 x 0.77 cm vascular, hyperechoic nodule with hypoechoic halo in the left lobe. No extrinsic masses are noted.

RIGHT LOBE: 3.39 x 1.88 x 1.70 cm
LEFT LOBE: 2.96 x 1.68 x 1.67 cm
ISTHMUS: 0.34 cm

There are slightly prominent lymph nodes measuring 1.76 x 1.41 x 0.75 cm and 1.85 x 1.19 x 0.71 cm in the right mid neck.

IMPRESSION

- 1. Heterogeneous bilateral thyroid lobes.**
- 2. 0.80 x 0.67 x 0.66 cm hypoechoic nodule with hyperechoic echoes and 0.94 x 0.28 cm hypoechoic nodule, right lobe.**
- 3. 0.87 x 0.87 x 0.63 cm hyperechoic nodule with calcification and 1.31 x 0.97 x 0.77 cm vascular, hyperechoic nodule with hypoechoic halo, left lobe.**
- 4. Slightly prominent lymph nodes measuring 1.76 x 1.41 x 0.75 cm and 1.85 x 1.19 x 0.71 cm, right mid neck.**

Correlation with clinical findings and appropriate follow-up suggested including 6-12 month follow-up study; consider MRI/CT.

Thank you for referring this patient.
Electronically signed by:



MELVIN S. FAIGUS, M.D.
Diplomate of the American Board of Radiology